

The HLA Class I Restricted CTL Response in HIV-1 Infection: Systematic Identification of Optimal Epitopes

Christian Brander and Bruce D. Walker

AIDS Research Center, Massachusetts General Hospital, 149 13th Street, Room 5234, Charlestown, MA 02129, USA, and Harvard Medical School Boston, MA USA

I. Introduction

Viral infection induces CD8+ cytotoxic T lymphocytes (CTL) responses by presenting viral antigen on MHC class I molecules at the surface of infected cells (1). A vigorous CTL response is also associated with Human Immunodeficiency Virus (HIV) infection (2). HIV specific CTL are thought to play an important role in the control of the virus during clinical latency and may influence the course of disease development (3-6). Furthermore, experiments in HIV-exposed, uninfected individuals suggest that CTL could provide protection from infection (7-9). The characterization of the antigenic epitopes involved in CTL induction is important not only for a better understanding of disease pathogenesis, but also for possible vaccine development. Since the first description of HIV specific CTL, the efforts of many laboratories have led to the identification of a large number of epitopes involved in the HIV-1 specific CTL response in infected persons (10, 11). A summary of those sequences is accessible in this database. It contains more than 200 peptides, which contain putative HLA class I restricted epitopes. In many cases, these peptides require further definition as to the minimal epitope or the restricting HLA molecule. Here, we have compiled a list of peptides that have been defined in terms of the minimal length that mediates the most effective target cell sensitization in the context of the restricting class I molecule. More than 60 peptides fulfilling these criteria were included in the last database update (11). Within the last 12 months, more than 30 new optimal epitopes have been identified, many of which are not yet published or are in submitted reports.

II. Presentation and recognition of viral antigen on MHC class I

Recent advances in the understanding of MHC class I antigen processing have provided insights into peptide selection for CTL responses. Viral antigen is presented to specific T cells as short peptides in association with MHC molecules (1, 12). These antigenic epitopes are derived from viral cytosolic proteins which undergo proteasome mediated processing (3, 13). A number of parameters likely influence peptide processing, including amino acids both within and outside the CTL epitope in the cytosol (14). Following processing, the peptides are translocated into the lumen of the ER by the TAP1/TAP2 heterodimer and bound to nascent MHC class I molecules. Both interactions (TAP and MHC) seem to have restrictive specificities and to allow only some peptides to be presented (3, 13, 15). Whereas peptide binding to MHC molecules has been extensively studied, both the impact of different TAP-alleles on the generation of peptides and the link between TAP alleles and HLA alleles are unclear (15-17). As revealed by peptide titration studies and peptide elution from the isolated MHC class I molecule, HLA class I restricted epitopes normally are about 8-12 amino acids in length and have certain anchor residues to bind into specific pockets of the MHC class I molecule. These pockets, designated A, B, C, D, E and F, are located in the antigen binding site on the class I molecule formed by two alpha helices, with the floor of the groove formed by a beta-sheet structure (18). As the pockets are located at the edge of the peptide binding groove, the anchor positions in the peptides are often in positions 1, 2 and 9 of the processed peptide (19). However, other residues can influence the peptide binding as well (20). Eluting peptides from various MHC class I molecules has led to the definition of MHC-allele-specific binding motifs (19-21). These motifs can be used to screen protein sequences in order to find potentially antigenic epitopes presented by a certain HLA allele (22). However, this approach only reflects peptide binding but not necessarily their involvement in the natural CTL response *in vivo*, as such evaluations do not analyze the effects of protein processing and transportation. Very

recently, the x-ray crystal structure and orientation of a T cell receptor (TCR) bound to a MHC/peptide complex was described (23). The V region of the α and the β chains, resembling antibody structures, contacts the presented peptide in its core region through the third complementarity-determining regions (CDR 3), forming a deep hydrophobic cavity. The CDR1 and CDR2 regions of the α -chain appear to contact the MHC/peptide complex around the N-terminal end of the embedded peptide, whereas the β chain CDR1 and CDR2 contact the C-terminus of the peptide. These findings may help to explain the biased use of certain V- β chain segments observed in various immune responses, i.e. influenza infection and different autoimmune diseases (24, 25).

III. Identification of HIV-1 derived CTL epitopes

Various methods have been used to identify HIV-1-derived, HLA class I restricted epitopes. The majority of epitopes have been defined using CTL lines or clones recognizing recombinant viral antigen, such as target cells infected with vaccinia virus constructs expressing selected HIV-1 proteins. The minimal epitope for these responses is determined by using synthetic peptides, followed by titration assays with truncated peptides to identify the optimal epitope (26). Alternatively, binding-motif based searches for potential epitopes has led to the identification of some optimal epitopes (22). However, to prove that the binding peptide is involved in the natural immune response to HIV, recognition of the naturally processed peptide by CTL in infected persons is required. Finally, elution of peptide from HLA class I molecules allows for the direct isolation of processed, class I binding epitopes (27). However, these epitopes are not necessarily immunogenic *in vivo* and may thus not induce a CTL response in infected humans.

In published studies, different methods have been used to identify optimal epitopes. In some cases the HLA restriction has not been shown or the optimal epitope has not been defined by titration assays using truncated peptides. The Los Alamos HIV Molecular Immunology Database contains all reported sequences with CTL activity, regardless of their definition as optimal epitopes. However, for many different questions and approaches such as the design of a epitope based vaccine, the knowledge of the optimal epitope is a prerequisite. In the list presented here (Table 1), we include only those peptides which fit rigid criteria: 1) the CTL recognize the naturally processed epitope, 2) titration assays with truncated peptides have identified the optimal epitope and 3) the restricting HLA molecule has been defined. A number of likely optimal epitopes were not included as these criteria were not met. For example, we did not include putative HLA-Cw8 restricted epitopes, since Cw8 is in a strong linkage disequilibrium with HLA-B14 and it is thus difficult to identify the restricting HLA allele by using (mis)matched cell lines (28, 29). However, these sequences can be found in other parts of the database.

Table 1 Best Defined HIV CTL Epitopes

HLA	Protein	AA	Isolate	Sequence	Reference	Ref No*
HLA-A2				2 C	Falk 91	21
				L V		
				I L		
	p17	77-85	LAI	SLYNTVATL	Johnson 91, Parker 92,94	35,36,37
	RT	346-354	LAI	VIYQYMDDL	T. Harrer	74
	RT	476-484	LAI	ILKEPVHGV	Walker 89, Tsomides 91	38,39
	gp41	818-827	LAI	SLLNATDIAV	Dupuis 95	40
	nef	136-145	LAI	PLTFGWICYKL	Haas 96, B. Maier and B. Autran	41,PC
	nef	180-189	LAI	VLEWRFD SRL	Haas 96, B. Maier and B. Autran	41,PC
	nef	190-198	LAI	AFHHVAREL	Hadida 95, B. Autran	42,PC

Table 1 (cont.) Best Defined HIV CTL Epitopes

HLA	Protein	AA	Isolate	Sequence	Reference	Ref No*
HLA-A3.1				23 C	Di Brino 93	43
				IF K		
				L Y		
	p17	18–26	LAI	KIRLRPGGK	Harrer 96	44
	p17	20–28	LAI	RLRPGGKKK	B. Culmann, D. Lewinsohn, S. Riddell	PC
	p17	20–29	LAI	RLRPGGKKKY	B. Wilkes, D. Ruhl	PC
	RT	325–333	LAI	AIFQSSMTK	S. Threlkeld	S
	gp120	37–46	LAI	TVYYGVPVWK	Johnson 94 b	45
	gp41	775–785	LAI	RLRDLLLIIVTR	Takahashi 91	46
	nef	73–82	LAI	QVPLRPMTYK	Koenig 90, Culmann 91	47,48
HLA-A11				2 C	Zhang 93	49
				I K		
				L		
	p17	84–92	LAI	TLYCVHQRI	T. Harrer	PC
	p24	349–359	III-B	ACQGVGGPGGHK	Sipsas 96	29
	RT	325–333	LAI	AIFQSSMTK	Johnson 94 a, Zhang 93, S. Threlkeld	50,49,S
	RT	508–517	LAI	IYQEPFKNLK	B. Culmann	PC
	nef	73–82	LAI	PLRPMTYK	Culmann 91	48
	nef	84–92	LAI	AVDLSSHFLK	Culmann 91	48
HLA-A19						
A*7401	RT	71–79	Clade A/B/D	ITLWQRPLV	S. Rowland-Jones	PC
HLA-A24						
	p17	28–36	LAI	KYKLKHIVW	D. Lewinsohn	PC
	gp120	53–62	LAI	LFCASDAKAY	Lieberman 92, Shankar 95	51,52
	gp41	591–598	LAI	YLKDQQLL	Dai 92	53
HLA-A25						
	p24	145–155	LAI	QAISPRTLNAW	I. Kurane, K. West	PC
	p24	203–212	LAI	ETINEEAAEW	Klenerman 96, C. Van Baalen 96	54,55
HLA-A26						
	p24	167–175	LAI	EVIPMFSAL	P. Goulder	IP-AIDS
	RT	593–603	LAI	ETFYVDGAANR	B. Wilkes, D. Ruhl	PC
HLA-A28						
A*6802	RT	71–79	Clade A/B/D	ITLWQRPLV	S. Rowland-Jones	PC
A*6802	RT	85–93	Clade D	DTVLEEMNL	S. Rowland-Jones	PC

Table 1 (cont.) Best Defined HIV CTL Epitopes

HLA	Protein	AA	Isolate	Sequence	Reference	Ref No*
HLA-A29						
	gp120	376–384	LAI	FNCGGGEFFY	C. Wilson	IP–JV
HLA-A31						
	gp41	775–785	LAI	RLRDLLLLIVTR	Safrit 94 a, Safrit 94 b	56,57
HLA-A32						
	RT	559–568	LAI	PIQKETWETW	Harrer 96	44
	gp120	419–427	HXB2	RIKQIINMW	Harrer 96	44
HLA-B7						
				123 C APR L	Engelhard 93	58
	p24	148–156	LAI	SPRTLNAWV	D. Lewinsohn	PC
	p24	179–187	LAI	ATPQDLNTM	B. Wilkes, D. Ruhl	PC
	gp120	303–312	LAI	RPNNNTRKSI	J. Safrit and R.A. Koup	PC
	gp41	843–851	LAI	IPRRIRQGL	B. Wilkes	PC
	nef	68–77	LAI	FPVTPQVPLR	Haas 96, B. Maier and B. Autran	41,PC
	nef	128–137	LAI	TPGPGVRYPL	Culmann 94, Haas 96, B. Maier and B. Autran	59,41,PC
HLA-B8						
				3 5 C K K I	Hill 92, Sutton 93	60,61
	p17	24–31	LAI	GGKKKYKL	Rowland-Jones 93, S. Reid	62,U
	p17	74–82	LAI	ELRSLYNTV	P. Goulder	S
	gp120	2–10	III-B	RVKEYQHL	Sipsas 96	29
	gp41	591–598	LAI	YLKDQQLL	Johnson 92, Shankar 95	63,52
	nef	13–20	LAI	WPTVRERM	P. Goulder	S
	nef	90–97	LAI	FLKEKGGL	Culmann 94, P. Goulder	59,PC
HLA-B14						
				23 56 C RL RI L KY HL F	Di Brino 94	64
	p24	183–191	LAI	DLNTMLNTV	Nixon 88, Johnson 92	65,63
	p24	298–306	LAI	DRFYKTLRA	Harrer 96	44
	gp41	589–597	LAI	ERYLKDQQL	Johnson 92	63
HLA-B15						
	gp120	375–383	LAI	SFNCGGEFF	C. Wilson	73
HLA-B18						
	nef	135–143	LAI	YPLTFGWCY	Culmann 91, Culmann 94	48,59

Table 1 (cont.) Best Defined HIV CTL Epitopes

HLA	Protein	AA	Isolate	Sequence	Reference	Ref No*
HLA-B27				2 C R K R	Jardetzky 91	66
	p17	18–27	LAI	KIRLRPGGKK	D. Lewinsohn	PC
	p17	19–27	LAI	IRLRPGGKK	D. Lewinsohn	PC
	p24	263–272	LAI	KRWIILGLNK	Nixon 88 b, Buseyne 93	67,68
	gp41	590–597	LAI	RYLKDQQL	Shankar 95	52
	gp41	791–799	LAI	GRRGWEALKY	Liebermann 92, J. Liebermann	51,PC
	nef	73–82	LAI	QVPLRPMTYK	B. Culmann	PC
	nef	105–114	LAI	RRQDILDLWI	P. Goulder	S
	nef	134–141	LAI	RYPLTFGW	B. Culmann	PC
	gag	260–269	HIV-2	RRWIQLGLQK	S. Rowland-Jones	PC
HLA-B35				2 C P Y S	Hill 92	60
	p17	36–44	LAI	WASRELERF	P. Goulder	S
	p17	124–132	LAI	NSSKVSQNY	Rowland-Jones 95	69
	p24	254–262	LAI	PPIPVGDIY	Rowland-Jones 95	69
	RT	262–270	LAI	TVLDVGDAY	B. Wilkes, D. Ruhl	PC
	RT	273–282	III-B	VPLDEDFRKY	Sipsas 96, Shiga 96	29,34
	RT	328–336	III-B	NPDIVIYQY	Sipsas 96, Shiga 96	29,34
	RT	342–350	LAI	HPDIVIYQY	Rowland-Jones 95	69
	gp120	42–52	LAI	VPVWKEATTTL	B. Wilkes, D. Ruhl	PC
	gp41	611–619	LAI	TAVPWNASW	Johnson 94 b	45
	nef	74–81	LAI	VPLRPMTY	Culmann 91, Culmann 94	48,59
	gag	245–253	HIV-2	NPVPVGNIY	Rowland-Jones 95	69
HLA-B37	nef	120–128	LAI	YFPDWQNYT	Culmann 91	48, PC
HLA-B39	p24	193–201	LAI	GHQAAMQML	I. Kurane, K. West	PC
HLA-B42	p17	20–29	LAI	RLRPGGKKKY	B. Wilkes, D. Ruhl	PC
	RT	438–446	LAI	YPGIKVRQL	B. Wilkes, D. Ruhl	PC
HLA-B45	RT	591–600	LAI	GAETFYVDGA	B. Wilkes, D. Ruhl	PC

Table 1 (cont.) Best Defined HIV CTL Epitopes

HLA	Protein	AA	Isolate	Sequence	Reference	Ref No*
HLA-B51						
	p24	325–333	LAI	NANPDCKTI	B. Wilkes, D. Ruhl	PC
	RT	295–302	III-B	TAFTTIPSI	Sipsas 96	29
	gp41	557–565	III-B	RAIEAQQHL	Sipsas 96	29
HLA-B52						
	p24	275–282	LAI	RMYSPTSI	B. Wilkes, D. Ruhl	PC
HLA-B53						
				2 C		60
				P Y	Hill 92	
				F		
				W		
	HIV-2 gag	173–181	HIV-2	TPYDINQML	Gotch 93	70
HLA-B55						
	gp120	42–51	LAI	VPVWKEATTT	Shankar 95	52
HLA-B57						
	p24	147–155	III-B	ISPRTLNAW	Johnson 91, P. Goulder	35,IP–ARHR
	p24	140–149	LAI	TSTLQEQIGW	P. Goulder	IP–ARHR
	p24	162–172	LAI	KAFSPEVIPMF	P. Goulder	IP–ARHR
	p24	240–249	LAI	TSTLQEQIGW	P. Goulder	PC
	p24	311–319	LAI	QASQEVKNW	P. Goulder	IP–ARHR
	p24	311–319	LAI	QASQDVKNW	P. Goulder	IP–ARHR
	nef	116–125	LAI	HTQGYFPDWQ	Culmann 91	48, PC
	nef	120–128	LAI	YFPDWQNYT	Culmann 91	48, PC
HLA-B58						
	p24	140–149	LAI	TSTLQEQIGW	P. Goulder	IP–ARHR
HLA-Bw62						
	p17	20–29	LAI	RLRPGGKKKY	Johnson 91, B. Wilkes, D. Ruhl	35,PC
	p24	268–277	LAI	LGLNKIVRMV	Johnson 91	35
	RT	415–426	III-B	LVGKLNWASQIY	P. Johnson	PC
	RT	476–485	LAI	ILKEPVHGVY	Johnson 91, P. Johnson	35,PC
	nef	84–91	LAI	AVDLSHFL	Culmann 94	59
	nef	117–127	LAI	TQGYFPDWQNY	B. Culmann	PC
HLA-Cw01.02						
	p24	168–175	LAI	VIPMFSAL	P. Goulder	S

Table 1 (cont.) Best Defined HIV CTL Epitopes

HLA	Protein	AA	Isolate	Sequence	Reference	Ref No*
HLA-Cw4				2 C	Falk 94	71
				Y F		
				P L		
				F M		
	gp120	380–388	LAI	SFNCGGEFF	Johnson 93, C. Wilson	72,73

*In Ref No column the following abbreviations apply:

PC = personal communication

S = submitted

IP-AIDS = in press - *AIDS*

IP-JV = in press - *Journal of Virology*

IP-ARHR = in press - *AIDS Research and Human Retroviruses*

IV. Analysis of HIV-1 derived CTL epitopes

The list of HIV derived CTL epitopes will continue to grow, as new epitopes are found and longer sequences are mapped to the optimal epitope. The list given in Table 1 should provide investigators with a reliable compilation of optimal CTL epitopes in HIV. With the growing number of defined optimal epitopes, some interesting characteristics are beginning to emerge. More epitopes which overlap with previously described ones have been identified, as have epitopes which can be restricted by multiple HLA alleles. These latter epitopes could be interesting candidates for peptide based vaccine development. This is the case for RT protein derived epitope AIFQSSMTK (a.a.325- a.a.333) which binds to HLA-A3 and HLA-A11, both members of the same binding motif superfamily (30). The gag-p17 derived peptide RLRPGGKKKY (a.a. 20-29) was found to be presented by at least three different HLA alleles (HLA-A3, -B42, -Bw62). Additional examples of epitopes binding to multiple HLA molecules are included in this list and can be conveniently found by searching the Los Alamos Database. Furthermore, it becomes clear that some areas contain multiple overlapping peptides. By minute alteration (i.e. truncation by or substitution of one amino acid), those epitopes may be able to bind to additional HLA molecules without losing the binding affinity to the original HLA molecule(s) nor escaping recognition by CTL specific for the original peptide. Different studies have shown that there can be wide cross recognition of variants of certain epitopes (8, 31, 32). This would suggest that a less conserved, but highly immunodominant epitope may serve as a superior immunogen for vaccine development than one which is highly conserved, but only rarely recognized. However, one must consider the potential adverse effects of sequence variation within, or flanking, these epitopes which may alter peptide processing and presentation. In general, it seems that the induction of a strong CTL response requires epitopes that bind tightly to the class I molecule. It has been suggested that the immunogenicity of potential CTL epitopes can be predicted more accurately by the dissociation rate than by the binding affinity (33). Although Table 1 lists numerous epitopes, little is known about their potential immunodominance, which may be especially important with regard to the development of peptide based vaccines (32, 34). The mechanisms leading to such immunodominance are not clear, although several reasons have been discussed. These include the preferential processing of certain epitopes by the proteasome due to favored flanking regions, the binding affinity to the class I molecule, the peptide selectivity of the TAP dimer, and a lack in the T cell repertoire. Thus, epitopes selected as peptide vaccine candidates may need to be tested not only for variability and binding affinities but rather for parameters like dissociation rate and immunodominance.

Acknowledgments

We would like to thank all the investigators who contributed unpublished data for inclusion in this list and in the Los Alamos HIV Molecular Immunology Database. This work was supported by the Schweizerische Stiftung fuer Medizinisch Biologische Stipendien, the Swiss National Foundation, NIH grants A128568 and A130914, and by a grant from the Pediatric AIDS Foundation (ARIEL Project for the prevention of HIV-transmission from mother to infant).

References

- [1] Oldstone MBA. (1989). Viral persistence. *Cell* **56**:517-520.
- [2] McMichael AJ and Walker BD. (1994). Cytotoxic T lymphocyte epitopes: Implications for HIV vaccines. *AIDS* **8** (suppl 1):S155-S173.
- [3] York IA and Rock KL. (1996). Antigen processing and presentation by the class I major histocompatibility complex. *Ann. Rev. Immunol.* **14**:369-396.
- [4] Koup RA, Safrit JT, Cao Y, Andrews CA, McLeod G, Borkowsky W Farthing C. and Ho DD. (1994). Temporal association of cellular immune responses with the initial control of viremia in primary human immunodeficiency virus type 1 syndrome. *J. Virol.* **68**:4650-4655.
- [5] Klein MR and Van Baalen CA. (1995). Kinetics of Gag-specific cytotoxic T lymphocyte responses during the clinical course of HIV-1 infection: A longitudinal analysis of rapid progressors and long-term asymptomatics. *J. Exp. Med.* **181**:1356-1372.
- [6] Harrer T, Harrer E, Kalams SA, Elbeik T, Staprans SI, Feinberg MB, Cao Y, Ho DD, Yilma T, Caliendo AM, Johnson RP, Buchbinder SP and Walker BD. (1995). Strong cytotoxic T cell and weak neutralizing antibody responses in a subset of persons with stable nonprogressing HIV type 1 infection. *AIDS Res. Human Retrov.* **12**:585-592.
- [7] Rowland-Jones SL, Nixon DF, Aldhous MC, Gotch F, Ariyoshi K, Hallam N, Kroll JS, Froebel K and McMichael A. (1993). HIV-specific cytotoxic T-cell activity in an HIV-exposed but uninfected infant. *Lancet* **341**:860-861.
- [8] Rowland-Jones S, Sutton J, Ariyoshi K, Dong T, Gotch F, McAdam S, Whitby D, Sabally S, Gallimore A, Corrah T, Takiguchi M, Schultz T, McMichael A and Whittle H. (1995). HIV-specific cytotoxic T-cells in HIV-exposed but uninfected Gambian women. *Nature Medicine* **1**:59-64.
- [9] Pinto LA, Sullivan J, Berzofsky JA, Clerici M, Kessler HA, Landay AL and Shearer GM. (1995). Env-specific cytotoxic T lymphocyte responses in HIV seronegative health care workers occupationally exposed to HIV contaminated body fluids. *J. Clin. Invest.* **96**:867-876.
- [10] Walker BD, Chakrabarti S, Moss B, Paradis TJ, Flynn T, Durno AG, Blumberg RS, Kaplan JC, Hirsch MS and Schooley RT. (1987). HIV-specific cytotoxic T lymphocytes in seropositive individuals. *Nature* **328**:345-8.
- [11] Brander C and Walker BD. (1995). The HLA class I restricted CTL response in HIV infection: identification of optimal epitopes. From: *HIV Molecular Immunology Database 1995*. Editors: Korber BTM, Brander C, Walker BD, Koup R, Moore JP, Haynes BF and Myers G. Publisher: Los Alamos National Laboratory: Theoretical Biology and Biophysics, Los Alamos, NM, USA. Part IV:1-9.
- [12] Goldberg AL and Rock KL. (1992). Proteolysis, proteasomes and antigen presentation. *Nature* **357**:375-9.
- [13] Rammensee HG, Falk K and Roetzschke O. (1993). MHC molecules as peptide receptors. *Curr. Opin. Immunol.* **5**:35-44.
- [14] Bergmann CC, Yao Q, Ho C and Buckwold SL. (1996). Flanking residues alter antigenicity and immunogenicity of multi-unit CTL epitopes. *J. Immunol.* **157**:3242-3249.
- [15] Heemels M and Ploegh HL. (1994). Substrate specificity of allelic variants of the TAP peptide transporter. *Immunity* **1**:775-784.
- [16] Shepherd JC, Schumacher TNM, Ashton-Rickardt PG, Imaeda S, Ploegh HL, Janeway CA and Tonegawa S. (1993). TAP 1-dependent peptide translocation *in vitro* is ATP dependent and peptide selective. *Cell* **74**:577-584.
- [17] Obst R, Armandola EA, Nijenhuis M, Momburg F and Haemmerling GJ. (1995). TAP polymorphism does not influence transport of peptide variants in mice and humans. *Eur. J. Immunol.* **25**:2170.
- [18] Bjorkman P, Saper M, Samraoui B, Bennett W, Strominger J and Wiley D. (1987). Structure of human class I histocompatibility antigen, HLA-A2. *Nature* **329**:506-511.
- [19] Falk K, Rotzschke O, Deres K, Metzger J, Jung G and Rammensee HG. (1991). Identification of naturally processed viral nonapeptides allows their quantification in infected cells and suggests an allele-specific T cell epitope forecast. *J. Exp. Med.* **174**:425-34.
- [20] Ruppert J, Sidney J, Celis E, Kubo RT, Grey HM and Sette A. (1993). Prominent role of secondary anchor residues in peptide binding to HLA-A2.1 molecules. *Cell* **74**:929-937.
- [21] Falk K, Rotzschke O, Stevanovic S, Jung G and Rammensee HG. (1991). Allele-specific motifs revealed by sequencing of self-peptides eluted from MHC molecules. *Nature* **351**:290-6.

- [22] Hobohm U and Meyerhans A. (1993). A pattern search method for putative anchor residues in T cell epitopes. *Eur. J. Immunol.* **23**:1271-1275.
- [23] Garcia KC, Degano M, Stanfield RL, Brunmark A, Jackson MR, Peterson PA, Teyton L and Wilson IA. (1996). An a/b T cell receptor structure at 2.5 Å and its orientation in the TCR-MHC complex. *Science* **274**:209-219.
- [24] Lehner PJ, Wang EC, Moss PAH, Williams S, Platt K, Friedman SM, Bell JI and Borysiewicz LK. (1995). Human HLA-A0201-restricted cytotoxic T lymphocyte recognition of Influenza A is dominated by T cells bearing the Vb17 gene segment. *J. Exp. Med.* **181**:79-91.
- [25] Moss PA, Rosenberg WM and Bell JI. (1992). The human T cell receptor in health and disease. *Ann. Rev. Immunol.* **10**:71-96.
- [26] Gnann JWW, Nelson JA and Oldstone MB. (1987). Fine mapping of an immunodominant domain in the transmembrane glycoprotein of human immunodeficiency virus. *J. Virol.* **61**:2639-41.
- [27] Tsomides TJ, Aldovini A, Johnson RP, Walker BD, Young RA and Eisen HN. (1994). Naturally processed viral peptides recognized by cytotoxic T lymphocytes on cells chronically infected by human immunodeficiency virus type 1 (HIV-1). *J. Exp. Med.* **180**:1283-1293.
- [28] Trowsdale J, Ragoussis J and Campbell RD. (1991). Map of the human MHC. *Immunology Today* **12**:443-446.
- [29] Sipsas NV, Kalams SA, Trocha A, He S, Blattner WA, Walker BD and Johnson RP. (1996). Identification of type-specific cytotoxic T lymphocyte responses to homologous viral proteins in laboratory workers accidentally infected with the human immunodeficiency virus-1. *J. Clin. Invest.* **In press**.
- [30] Threlkeld SC, Wentworth P, Kalams S, Johnson P, Wilkens BM, Ruhle DJ, Sidney J, Walker BD and Sette A. (1996). Degenerate and promiscuous recognition by cytotoxic T lymphocyte (CTL) of peptides presented by the MHC class I A3-like superfamily: Implications for vaccine development. **Submitted**.
- [31] Safrit J, Andrews CA, Tuofu Z, Ho DD and Koup RA. (1994). Characterization of human immunodeficiency virus type 1-specific cytotoxic T lymphocyte clones isolated during acute seroconversion: Recognition of autologous virus sequences within a conserved immunodominant epitope. *J. Exp. Med.* **179**:463-472.
- [32] Brander C, Hartman KE, Trocha AK, Jones ND, He S, Wentworth P, Hermanson G, Buchbinder SP, Johnson RP, Walker BD and Kalams SA. (1996). Identification of the immunodominant HLA-A*02.01 restricted CTL responses in HIV infection: strong CTL responses contrast with only rare occurrence of potential viral escape variants. **Submitted**.
- [33] Van der Burg S, Visseren MJ, Brandt RM, Kast WM and Melief CJ. (1996). Immunogenicity of peptides bound to MHC class I molecules depends on the MHC peptide complex stability. *J. Immunol.* **156**:3308-3314.
- [34] Shiga H, Shioda T, Tomiyama H, Takamiya Y, Oka S, Kimura S, Yamaguchi Y, Gojoubori T, Rammensee HG, Miwa K and Takiguchi M. (1996). Identification of multiple HIV-1 cytotoxic T-cell epitopes presented by human leukocyte antigen B35 molecule. *AIDS* **10**:1075-1083.
- [35] Johnson RP, Trocha A, Yang L, Mazzara GP, Panicali DL, Buchanan TM and Walker BD. (1991). HIV-1 gag-specific cytotoxic T lymphocytes recognize multiple highly conserved epitopes. Fine specificity of the gag-specific response defined by using unstimulated peripheral blood mononuclear cells and cloned effector cells. *J. Immunol.* **147**:1512-1521.
- [36] Parker KC, Bednarek MA, Hull LK, Utz U, Cunningham B, Zweerink HJ, Biddison WE and Coligan JE. (1992). Sequence motifs important for peptide binding to the human MHC class I molecule, HLA-A2. *J. Immunol.* **149**:3580-3587.
- [37] Parker KC, Bednarek MA and Coligan JE. (1994). Scheme for ranking potential HLA-A2 binding peptides based on independent binding of individual peptide side-chains. *J. Immunol.* **152**:163-175.
- [38] Walker BD, Flexner C, Birch-Limberger K, Fisher L, Paradis TJ, Aldovini A, Young R, Moss B and Schooley RT. (1989). Long-term culture and fine specificity of human cytotoxic T-lymphocyte clones reactive with human immunodeficiency virus type 1. *Proc. Natl. Acad. Sci. USA*. **86**:9514-9518.
- [39] Tsomides TJ, Walker BD and Eisen HN. (1991). An optimal viral peptide recognized by CD8+ T cells binds very tightly to the restricting class I major histocompatibility complex protein on intact cells but not to the purified class I protein. *Proc. Natl. Acad. Sci. USA*. **88**:11276-11280.
- [40] Dupuis M, Kundu SK and Merigan TC. (1995). Characterization of HLA-A*0201-restricted cytotoxic T cell epitopes in conserved regions of the HIV type 1 gp160 protein. *J. Immunol.* **155**:2232-2239.
- [41] Haas G, Plikat U, Debre P, Lucchiari M, Katlama C, Dudoit Y, Bonduelle O, Bauer M, Ihlenfeldt H, Jung G, Maier B, Meyerhans A and Autran B. (1996). Dynamics of viral variants in HIV-1 Nef and specific cytotoxic T lymphocytes *in vivo*. *J. Immunol.* **157**:4212-4221.

- [42] Hadida F, Haas G, Zimmermann G, Hosmalin A, Spohn R, Samri A, Jung G, Debre P and Autran B. (1995). CTLs from lymphoid organs recognize an optimal HLA-A2 restricted and HLA-B52-restricted nonapeptide and several epitopes in the C-terminal region of HIV-1 Nef. *J. Immunol.* **154**:4174–4186.
- [43] DiBrino M, Parker KC, Shiloach J, Knierman M, Lukszo J, Turner RV, Biddison WE and Coligan JE. (1993). Endogenous peptides bound to HLA-A3 possess a specific combination of anchor residues that permit identification of potential antigenic peptides. *Proc. Natl. Acad. Sci. USA.* **90**:1508–1512.
- [44] Harrer T, Harrer E, Kalams SA, Barbosa P, Trocha A, Johnson RP, Elbeik T, Feinberg MB, Buchbinder SP and Walker BD. (1996). Cytotoxic T lymphocytes in asymptomatic long-term nonprogressing HIV-1 infection. Breadth and specificity of the response and relation to *in vivo* viral quasiespecies in a person with prolonged infection and low viral load. *J. Immunol.* **156**:2616–2623.
- [45] Johnson RP, Hammond SA, Trocha A, Siliciano RF and Walker BD. (1994). Induction of a major histocompatibility complex class I-restricted cytotoxic T lymphocyte response to a highly conserved region of human immunodeficiency virus type 1 (HIV-1) gp120 in seronegative humans immunized with a candidate HIV-1 vaccine. *J. Virol.* **68**:3145–3153.
- [46] Takahashi K, Dai LC, Fuerst TR, Biddison WE, Earl PL, Moss B and Ennis FA. (1991). Specific lysis of human immunodeficiency virus type 1-infected cells by a HLA-A3.1-restricted CD8+ cytotoxic T-lymphocyte clone that recognizes a conserved peptide sequence within the gp41 subunit of the envelope protein. *Proc. Natl. Acad. Sci. USA.* **88**:10277–10281.
- [47] Koenig S, Fuerst TR, Wood LV, Woods RM, Suzich JA, Jones GM, de la Cruz VF, Davey RT Jr, Venkatesan S, Moss B. et al. (1990). Mapping the fine specificity of a cytolytic T cell response to HIV-1 nef protein. *J. Immunol.* **145**:127–135.
- [48] Culmann B, Gomard E, Kieny MP, Guy B, Dreyfus F, Saimot AG, Sereni D, Sicard D and Levy JP. (1991). Six epitopes reacting with human cytotoxic CD8+ T cells in the central region of the HIV-1 Nef protein. *J. Immunol.* **146**:1560–2.
- [49] Zhang QJ, Gavioli R, Klein G and Masucci MG. (1993). An HLA-A11-specific motif in nonamer peptides derived from viral and cellular proteins. *Proc. Natl. Acad. Sci. USA.* **90**:2217–2221.
- [50] Johnson RP and Walker BD. (1994). CTL in HIV-1 infection: Responses to structural proteins. *J. Curr. Topics. Microbiol. Immunol.* **189**:35–63.
- [51] Lieberman J, Fabry JA, Kuo MC, Earl F, Moss B and Skolnik PR. (1992). Cytotoxic T lymphocytes from HIV-1 seropositive individuals recognize immunodominant epitopes in gp160 and reverse transcriptase. *J. Immunol.* **148**:2738–2747.
- [52] Shankar P, Fabry JA, Fong D and Lieberman J. (1995). Three regions of gp160 contain overlapping CTL epitopes restricted by multiple HLA class I alleles. (abstract) *J. Cellular. Biochem.* S21B: D4–345.
- [53] Dai LC, West K, Littau R, Takahashi K and Ennis FA. (1992). Mutation of human immunodeficiency virus type 1 at amino acid 585 on gp41 results in loss of killing by CD8+ A24-restricted cytotoxic T lymphocytes. *J. Virol.* **66**:3151–3154.
- [54] Klenerman P, Luzzi G, McIntyre K, Phillips R and McMichael A. (1996). Identification of a novel HLA-A25 restricted epitope in a conserved region of p24 gag (positions 71–80). *AIDS.* **10**:348–350.
- [55] Van Baalen CA, Klein MR, Huisman RC, Dings ME, Kerkhof Garde SR, Geretti AM, Gruters R, Van Els CA, Miedema F and Osterhaus AD. (1996). Fine-specificity of cytotoxic T lymphocytes which recognize conserved epitopes of the Gag protein of human immunodeficiency virus type 1. *J. Gen. Virol.* **77**:1659–1665.
- [56] Safrit JT, Lee AY, Andrews CA and Koup RA. (1994). A region of the third variable loop of HIV-1 gp120 is recognized by HLA-B7-restricted CTLs from two acute seroconversion patients. *J. Immunol.* **153**:3822–3830.
- [57] Safrit JT, Andrews CA, Zhu T, Ho DD and Koup RA. (1994). Characterization of human immunodeficiency virus type 1-specific cytotoxic T lymphocyte clones isolated during acute seroconversion: recognition of autologous virus sequences within a conserved immunodominant epitope. *J. Exp. Med.* **179**:463–472.
- [58] Engelhard VH, Huczko EL and Bodener W. (1993). Peptides bound to HLA-B7 determined by mass spectrometry (abstract) *J. Cell. Biochem. Suppl.* 17C:56.
- [59] Culmann-Penciolelli B, Lamhamedi-Cherradi S, Couillin I, Guegan N, Levy JP, Guillet JG and Gomard E. (1994). Identification of multirestricted immunodominant regions recognized by cytolytic T lymphocytes in the human immunodeficiency virus type 1 nef protein. *J. Virol.* **68**:7336–7343.
- [60] Hill AV, Elvin J, Willis AC, Aidoo M, Allsopp CE and Gotch FM. (1992). Molecular analysis of the association of HLA-B53 and resistance to severe malaria. *Nature.* **360**:434–439.

- [61] Sutton J, Rowland-Jones S, Rosenberg W, Nixon D, Gotch F, Gao XM, Murray N, Spoonas A, Driscoll P, Smith M, Willis A and McMichael AJ. (1993). A sequence pattern for peptides presented to cytotoxic T lymphocytes by HLA B8 revealed by analysis of epitopes and eluted peptides. *Eur. J. Immunol.* **23**:447–453.
- [62] Rowland-Jones SL, Powis SH, Sutton J, Mockridge I, Gotch FM, Murray N, Hill AB, Rosenberg WM, Trowsdale J and McMichael AJ. (1993). An antigen processing polymorphism revealed by HLA-B8-restricted cytotoxic T lymphocytes which does not correlate with TAP gene polymorphism. *Eur. J. Immunol.* **23**:1999–2004.
- [63] Johnson RP, Trocha A, Buchanan TM and Walker BD. (1992). Identification of overlapping HLA class I-restricted cytotoxic T cell epitopes in a conserved region of the human immunodeficiency virus type 1 envelope glycoprotein: definition of minimum epitopes and analysis of the effects of sequence variation. *J. Exp. Med.* **175**:961–971.
- [64] DiBrino M, Parker KC, Margulies DH, Shiloach J, Turner RV, Garfield M, Biddison WE and Coligan JE. (1994). The HLA-B14 peptide binding site can accommodate peptides with different combinations of anchor residues. *J. Biol. Chem.* **269**:32426–32434.
- [65] Nixon DF and McMichael AJ. (1988). Cytotoxic T cell recognition of HIV proteins and peptides. *AIDS.* **5**:1049–1059.
- [66] Jardetzky TS, Lane WS, Robinson RA, Madden DR and Wiley DC. (1991). Identification of self peptides bound to purified HLA-B27. *Nature.* **353**:326–329.
- [67] Nixon DF, Townsend AR, Elvin JG, Rizza CR, Gallwey J and McMichael A. (1988). HIV-1 gag specific cytotoxic T lymphocytes defined with recombinant vaccinia virus and synthetic peptides. *Nature.* **336**:484–487.
- [68] Buseyne F, McChesney M, Porrot F, Kovarik S, Guy B and Riviere Y. (1993). Gag-specific cytotoxic T lymphocytes from human immunodeficiency virus type 1-infected individuals: Gag epitopes are clustered in three regions of the p24 gag protein. *J. Virol.* **67**:694–702.
- [69] Rowland-Jones S, Sutton J, Ariyoshi K, et al. (1995). HIV-specific cytotoxic T-cells in HIV-exposed but uninfected Gambian women. *Nature Med.* **1**:59–64.
- [70] Gotch F, McAdam SN, Allsopp CE, Gallimore A, Elvin J, Kieny MP, Hill AV and McMichael AJ. (1993). Cytotoxic T cells in HIV-2 seropositive Gambians. Identification of a virus-specific MHC-restricted peptide epitope. *J. Immunol.* **151**:3361–3369.
- [71] Falk K, Rötzschke O and Grahovac B. (1994). Allele specific peptide motifs of HLA-C molecules. *Proc. Natl. Acad. Sci. USA.* **90**:12005–12009.
- [72] Johnson RP, Trocha A, Buchanan TM and Walker BD. (1993). Recognition of a highly conserved region of human immunodeficiency virus type 1 gp120 by an HLA-Cw4 restricted cytotoxic T lymphocyte clone. *J. Virol.* **67**:438–445.
- [73] Wilson CC, Kalams SA, Wilkes BA, Ruhl DH, Gao F, Hahn BH, Hanson IC, Luzuriaga K, Wolinsky S, Koup R, Buchbinder SP, Johnson RP, and Walker BD. (1996). Overlapping Epitopes in Human Immunodeficiency Virus Type 1 gp120 presented by HLA A, B and C Molecules: Effects of viral variation on Cytotoxic T-Lymphocyte Recognition. *J. Virol.* **71**:1256–1264.
- [74] Harrer E, Harrer T, Barbosa P, Feinberg M, Johnson RP, Buchbinder S and Walker BD (1996). Recognition of Highly Conserved YMDD Region in the Human Immunodeficiency Virus Type 1 Reverse Transcriptase by HLA-A2-Restricted Cytotoxic T Lymphocyte from an Asymptomatic Long-Term NonProgressor. *J. of Infectious Diseases*, **173**(2): 476–9.